

# BIOMIN<sup>®</sup> CALCIUM POWDER

## 5-0-0



## Chelated Minerals for Soil and Foliar Application

### PRODUCT INFORMATION

**BIOMIN CALCIUM** contains nitrogen and chelated calcium for soil and/or foliar applications. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. **BIOMIN CALCIUM** is designed to prevent and correct calcium deficiency and to boost crops during critical or fast growing periods. **BIOMIN CALCIUM** may be applied with nitrogen fertilizers.

**BIOMIN CALCIUM** can be applied to ALL commercial crops including all vegetables, Potatoes, Carrots, Onions, all berries and strawberries, all Brassicas, all fruit trees, Avocados, Persimmons, Citrus, all Nut trees, Vineyards, Kiwifruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Ginger, Ornamentals, Cut flowers and Pastures.

### DIRECTIONS AND GENERAL RECOMMENDATIONS: SOIL APPLICATION

**BIOMIN CALCIUM** may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1.5-5.0 pounds per acre (1.7-5.6 Kg / Ha) depending on crop, soil conditions and application methods. For maintenance, 1.5 pounds per acre (1.7 Kg / Ha) are usually recommended.

### FOLIAR APPLICATION

**FIELD CROPS:** Apply 1.0-1.5 pounds of **BIOMIN CALCIUM** per acre (1.1-1.7 Kg/Ha) when crops are 6-12 inches (15-30 cm) high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

**VEGETABLES and POTATOES:** Apply 1.0-1.5 pounds of **BIOMIN CALCIUM** per acre (1.1-1.7 Kg/Ha) at about 2-4 weeks after emergence or if

### GUARANTEED ANALYSIS

Total Nitrogen (N).....5.0%  
5% Water Soluble Organic Nitrogen  
Calcium (Ca).....15.0%  
15% Chelated Calcium

Derived from Calcium Carbonate and Citric Acid.  
Nitrogen derived from Glycine

### PREVENTS AND CORRECTS CALCIUM DEFICIENCY

### KEEP OUT OF REACH OF CHILDREN

Manufactured by:  
JH BIOTECH, INC.  
P.O. Box 3538  
Ventura, CA 93006  
U.S.A.

### NET CONTENTS

- 2.5 pounds (1,1 Kg)
- 5 pounds (2,3 Kg)
- 25 pounds (11,4 Kg)
- 55 pounds (25 Kg)

Lot No.:

U.S. Pat.# 5,504,055

deficiency symptoms appear. Repeat the application 2-3 weeks later.

**GREENHOUSE CROPS:** Apply 1.0-1.5 pounds of **BIOMIN CALCIUM** per acre (1.1-1.7 Kg/Ha) 2-3 weeks after emergence or transplanting, or when calcium deficiencies first appear. Repeat the application every one or two weeks.

**FRUIT TREES AND GRAPES:** Apply 1.0-1.5 pounds of **BIOMIN CALCIUM** per acre (1.1-1.7 Kg/Ha) in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

**BIOMIN CALCIUM** is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 200 times of water. For best results and more specific recommendations, contact your nearest distributor. Always perform a compatibility test before mixing any products with **BIOMIN CALCIUM**.

### LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapco.org/metals.htm>.

**JH Biotech, Inc.**

0401-D-A-2  
Copyright © JH Biotech, Inc.  
All rights reserved.