



# BIOMIN® COPPER

## 2-0-0



## Amino Acid Chelated Mineral for Soil and Foliar Application

### PRODUCT INFORMATION:

**BIOMIN COPPER** contains amino acid chelated minerals for soil and foliar application. It is completely available and non-phytoxic to plants when applied according to directions. It is designed to prevent and correct mineral deficiency and to boost crops during critical stage or fast growing periods. It can be applied in combination with nitrogen fertilizer.

### DIRECTIONS AND

### GENERAL RECOMMENDATIONS:

#### SHAKE WELL BEFORE USING

**SOIL APPLICATION:** **BIOMIN COPPER** can be applied before or at planting as banding or broadcast. The application rate is 1 quart to 1 gallon per acre (2.5 to 10 liters/hectare) depending on soil conditions and application methods. For maintenance, 1 quart per acre (2.5 liters/hectare) is recommended. **BIOMIN COPPER** should be diluted with enough water to cover the entire acre.

### FOLIAR APPLICATION:

**FIELD CROPS:** Apply 1-4 pints per acre (1.5 to 5 liters/hectare) of **BIOMIN COPPER** when crops are about 6-12 inches (15 - 30 cm) high or when deficiency symptoms appear. Repeat the application in 2 - 4 weeks if necessary.

**FRUIT TREES AND GRAPES:** Apply 1-4 pints per acre (1.5 to 5 liters/hectare) starting at bud break or when deficiency symptoms appear.

### GUARANTEED ANALYSIS

Total Nitrogen (N) .....2.0%  
2.0% Water Soluble Organic Nitrogen

Copper (Cu).....4.0%  
4.0% Chelated Copper

Contains Copper Sulfate, Citric Acid and Glycine as chelating agents.  
(Nitrogen derived from Glycine.)

### PREVENT AND CORRECT MINERAL DEFICIENCIES

### KEEP OUT OF REACH OF CHILDREN

**Manufactured by:**  
JH BIOTECH, INC.  
P.O. Box 3538  
Ventura, CA 93006

### NET CONTENTS:

- 1 Pint (0.47 liters)
- 2.5 Gallons (9.45 liters)
- 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters)
- 55 Gallons (207.9 liters)
- Other: \_\_\_\_\_ Gallons

**Weight per Gallon:** 10.8 lbs. (4.9 kg.)

**U.S. Pat.# 5,504,055**

Lot No.:

Repeat the application in about 4 weeks if necessary.

**VEGETABLE CROPS:** Apply 1-4 pints per acre (1.5 to 5 liters/hectare) of **BIOMIN COPPER** at about 2 - 4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1 - 3 weeks if necessary.

For best results, apply **BIOMIN COPPER** when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, it should be diluted with at least 20 times of water. **BIOMIN COPPER** will lower high pH of water and decrease the hydrolysis of most pesticides.

**Biomim Copper Can Be Applied To** Strawberry, Lettuce, Tomato, Celery, Stone Fruits, Vegetables, Grapes, Commercial Turf, And Ornamental Gardens.

### LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>