

BIOMIN[®] COPPER POWDER

5-0-0

Chelated Minerals for Soil and Foliar Application



PRODUCT INFORMATION

BIOMIN COPPER contains nitrogen and chelated copper for soil and/or foliar applications. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. **BIOMIN COPPER** is designed to prevent and correct copper deficiency and to boost crops during critical or fast growing periods. **BIOMIN COPPER** may be applied with nitrogen fertilizers.

BIOMIN COPPER can be applied to ALL commercial crops including all vegetables, Potatoes, Carrots, Onions, all berries and strawberries, all Brassicas, all fruit trees, Avocados, Persimmons, Citrus, all Nut trees, Vineyards, Kiwifruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Ginger, Ornamentals, Cut flowers and Pastures.

DIRECTIONS AND

GENERAL RECOMMENDATIONS:

SOIL APPLICATION

BIOMIN COPPER may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1.5-5.0 pounds per acre (1.7-5.6 Kg/Ha) depending on crop, soil conditions and application methods. For maintenance, 1.5 pounds per acre (1.7 Kg/Ha) are usually recommended.

FOLIAR APPLICATION

FIELD CROPS: Apply 1.0-1.5 pounds of **BIOMIN COPPER** per acre (1.1 -1.7 Kg/Ha) when crops are 6-12 inches (15-30 cm) high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

VEGETABLES and POTATOES: Apply 1.0-1.5 pounds of **BIOMIN COPPER** per acre (1.1 -1.7 Kg/Ha) at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later.

GUARANTEED ANALYSIS

Total Nitrogen (N).....5.0%
5% Water Soluble Organic Nitrogen
Copper (Cu).....17.0%
17% Chelated Copper

Derived from Copper Sulfate and Citric Acid.
Nitrogen derived from Glycine

PREVENTS AND CORRECTS COPPER DEFICIENCY

KEEP OUT OF REACH OF CHILDREN

Manufactured by:

JH BIOTECH, INC.
P.O. Box 3538
Ventura, CA 93006
U.S.A.

NET CONTENTS

- 2.5 pounds (1.1 Kg)
- 5 pounds (2.3 Kg)
- 25 pounds (11.4 Kg)
- 55 pounds (25 Kg)

Lot No.:

U.S. Pat.# 5,504,055

GREENHOUSE CROPS: Apply 1.0-1.5 pounds of **BIOMIN COPPER** per acre (1.1-1.7 Kg/Ha) 2-3 weeks after emergence or transplanting, or when copper deficiencies first appear. Repeat the application every one or two weeks.

FRUIT TREES AND GRAPES: Apply 1.0-1.5 pounds of **BIOMIN COPPER** per acre (1.1-1.7 Kg/Ha) in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

BIOMIN COPPER is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 200 times of water. For best results and more specific recommendations, contact your nearest distributor. Always perform a compatibility test before mixing any products with **BIOMIN COPPER**.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

JH Biotech Inc.

0402-D-A-2
Copyright © JH Biotech, Inc.
All rights reserved.