



BIOMIN[®] IRON

2-0-0



Amino Acid Chelated Mineral for Soil and Foliar Application

PRODUCT INFORMATION:

BIOMIN IRON contains amino acid chelated mineral for soil and foliar applications. It is completely available and non-phytoxic to plants when applied according to directions. It is designed to prevent and correct mineral deficiency and to boost crops during critical stages or fast growing periods. It can be applied in combination with nitrogen fertilizer.

DIRECTIONS AND GENERAL

RECOMMENDATIONS:

SHAKE WELL BEFORE USING

SOIL APPLICATION: BIOMIN IRON can be applied before or at planting as banding or broadcast. The application rate is 1 quart to 1 gallon per acre depending on soil conditions and application methods. For maintenance, 1 quart per acre is recommended. **BIOMIN IRON** should be diluted with enough water to cover the entire acre.

FOLIAR APPLICATION:

FIELD CROPS:

Apply 1-4 pints per acre (1.5-5 liters/hectare) of **BIOMIN IRON** when crops are about 4-6 inches (15-30 cm) high or when deficiency symptoms appear. Repeat the application in 2 - 4 weeks if necessary.

FRUIT TREES AND GRAPES:

Apply 1-4 pints per acre (1.5-5 liters/hectare) starting at bud break or when deficiency

GUARANTEED ANALYSIS

Total Nitrogen (N)	2.0%
2.0% Water Soluble Organic Nitrogen	
Iron (Fe).....	5.0%
5.0% Chelated Iron	

Contains Ferrous Sulfate, Citric Acid and Glycine as chelating agents. (Nitrogen derived from Glycine.)

PREVENT AND CORRECT MINERAL DEFICIENCIES

KEEP OUT OF REACH OF CHILDREN

Manufactured by:

JH BIOTECH, INC.

P.O. Box 3538

Ventura, CA 93006

Net Contents:

- 1 Pint (0.47 liter)
- 2.5 Gallons (9.45 liters)
- 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters)
- 55 Gallons (207.9 liters)
- Other: _____ Gallons

Weight per Gallon: 10.5 lbs. (4.8 kg)

U.S. Pat.# 5,504,055

Lot No.:

symptoms appear. Repeat the application in about 4 weeks if necessary.

VEGETABLE CROPS:

Apply 1-4 pints per acre (1.5-5 liters/hectare) of **BIOMIN IRON** at about 2 - 4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1 - 3 weeks if necessary.

For best results, apply **BIOMIN IRON** when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, it should be diluted with at least 100 times of water. **BIOMIN IRON** will lower high pH of water and decrease the hydrolysis of most pesticides.

Biomin Iron Can Be Applied To

Strawberry, Lettuce, Tomato, Celery, Stone Fruit, Grapes, Vegetables, Commercial Turf, and Ornamental Gardens.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available online at <http://www.aapfco.org/metals.htm>

JH Biotech Inc.

0303-D-A-2
Copyright © 2009 JH Biotech, Inc.
All rights reserved.