



OMRI[™]
Listed

BIOMIN[®] ZINC

2-0-0



Amino Acid Chelated Mineral for Soil and Foliar Application

PRODUCT INFORMATION

BIOMIN ZINC contains amino acid chelated mineral for soil and/or foliar application. Completely bioavailable and non-phyto-toxic to plants when applied in accordance to directions. Designed to prevent and correct micronutrient deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

DIRECTIONS AND GENERAL

RECOMMENDATIONS:

SHAKE WELL BEFORE USING

SOIL APPLICATION: **BIOMIN ZINC** can be applied before or at planting as banding or broadcast. The application rate is 1 quart to 1 gallon per acre (2.5 to 10 liters/hectare) depending on soil conditions and application methods. For maintenance, 1 quart per acre (2.5 liters/hectare) is recommended. **BIOMIN ZINC** should be diluted with enough water to cover the entire acre.

FOLIAR APPLICATION:

FIELD CROPS: Apply 1-4 pints per acre (1.5-5 liters/hectare), when crops are about 4-6 inches (10 - 20 cm) high or when deficiency symptoms appear. Repeat the application in 2 - 4 weeks if necessary.

FRUIT TREES AND GRAPES: Apply 1-4 pints per acre (1.5-5 liters/hectare) starting at bud break or when deficiency

GUARANTEED ANALYSIS

Total Nitrogen(N).....2.0%
2.0% Water Soluble Organic Nitrogen
Zinc (Zn).....7.0%
7.0% Chelated Zinc

Derived from Zinc Sulfate, Citric Acid and Glycine. Nitrogen derived from Glycine.

PREVENT AND CORRECT MINERAL DEFICIENCIES

KEEP OUT OF REACH OF CHILDREN

Manufactured by:

JH BIOTECH, INC.
P.O. Box 3538
Ventura, CA 93006
U.S.A.

NET CONTENTS:

- 1 Pint (0.47 liters)
- 2.5 Gallons (9.45 liters)
- 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters)
- 55 Gallons (207.9 liters)
- Other: _____ Gallons

Weight per Gallon: 10.9 lbs. (4.9 kg)

U.S. Pat.# 5,504,055

symptoms appear. Repeat the application in about 4 weeks if necessary.

VEGETABLE CROPS: Apply 1-4 pints per acre (1.5-5 liters/hectare) at about 2 - 4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1-3 weeks if necessary.

For best results, apply when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, it should be diluted with at least 100 times of water. **BIOMIN ZINC** will lower high pH of water and decrease the hydrolysis of most pesticides.

BiomIn Zinc Can Be Applied To

Strawberry, Lettuce, Tomato, Celery, Grapes, Stone Fruits, Tree Fruits, Vegetables, Commercial Turf And Ornamental Gardens.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label. Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

Lot No.:

0306-D-A-1

JH Biotech, Inc.