

NUTRI-AID™

6-30-30

Water Soluble Fertilizer With Micronutrients



GUARANTEED ANALYSIS:

Total Nitrogen (N)	6%
6.0% Ammoniacal Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	30%
Soluble Potash (K ₂ O)	30%
Boron (B)	0.02%
Copper (Cu)	0.05%
0.05% Chelated Copper	
Iron (Fe)	0.10%
0.10% Chelated Iron	
Manganese (Mn)	0.05%
0.05% Chelated Manganese	
Zinc (Zn)	0.05%
0.05% Chelated Zinc	

Derived from ammonium phosphate, potassium sulfate, sodium borate, copper sulfate, ferrous sulfate, manganese sulfate and zinc sulfate and citric acid.

DIRECTIONS:

DILUTE APPLICATIONS:	3 to 5 lbs. per 100 gallons of water
SEMI-CONCENTRATE:	5 to 10 lbs. per acre in 40 to 100 gallons of water
AERIAL APPLICATION:	3 to 10 lbs. per acre in sufficient water for good coverage

Fill tank with water and slowly add **NUTRI-AID** while agitating. Continue to agitate until all **NUTRI-AID** is completely dissolved.

APPLICATION IN IRRIGATION WATER:

Solid Set, Overhead, Drip Sprinkler or Pivot Irrigation Systems: Dissolve **NUTRI-AID** in mix tank and apply 8-10 lbs. per acre in irrigation water. Use of anti-siphon or check valve is recommended to prevent solution from running back into primary water supply.

NUTRI-AID is generally compatible with pesticides and other fertilizers. Determine compatibility with a jar test before mixing products.

AVOCADOS: Use 10 to 20 lbs. per application in sufficient water to thoroughly wet foliage.

BEANS, PEAS, LENTILS, RAPESEED: (Canola) Use 5 to 10 lbs. per acre. Apply at first sign of bloom. Repeat 2 weeks later.

BERRIES: (Strawberries, raspberries, blackberries, blueberries, etc.) Use 5 to 10 lbs. per acre in spring on new growth. Repeat at pre-bloom and 3 weeks later. Use sufficient water for good coverage.

CARROTS, BEETS, SWEET POTATOES, AND OTHER VEGETABLE ROOT CROPS: Use 5 to 10 lbs. per acre per application.

CITRUS INCLUDING ORANGE, LEMON, GRAPEFRUIT, TANGERINE: Use 10 to 20 lbs. per acre per application.

COLE CROPS, LETTUCE, LEAFY VEGETABLES: Use 5 to 10 lbs. per acre. Apply right after thinning or transplanting. Repeat every 10 to 14 days.

CRANBERRIES: Use 5 to 10 lbs. per acre. Apply at popcorn stage. Repeat at hooking and 3 weeks later.

DECIDUOUS FRUITS: (Apples, pears, cherries, plums, prunes, peaches, almonds, walnuts, filberts, pistachios and pecans) Apply in delayed dormant, pre-bloom, and post-bloom sprays at 3 to 5 lbs. per 100 gal. in dilute sprays. Apply 10 to 15 lbs. per acre in concentrate sprays. Avoid combining with material that will cause russetting.

FIELD CROPS: (Mint, alfalfa, clover) Use 5 to 10 lbs. per acre. Apply when plants have 6" to 8" new growth. Repeat application to new growth after each cutting.

GRAPES: Ground sprayers: Use 5 to 10 lbs. per acre in 50 to 100 gal. of water. Air Application: Use 5 to 8 lbs. per acre in 10 gal. of water. Apply when leaves emerge, pre-bloom and post-bloom until grapes are 1/4 size.

HOPS: Ground sprayers: Use 5 to 10 lbs. per acre in 100-200 gallons of water. Air Application: Use 5 to 8 lbs. per acre in no less than 10 gal. of water. Apply when hops are halfway up wire and again when they reach the top of the wire.

MELONS, CUCUMBERS AND OTHER CUCURBITS: Use 5 to 10 lbs. per acre per application. Apply two or more applications every 2 to 3 weeks beginning with pre-bloom.

MINT: Use 5 to 10 lbs. per acre per application.

ONIONS (GREEN AND DRY), GARLIC: Use 5 to 10 lbs. per acre per application.

POTATOES: Use 5 to 10 lbs. per acre. Apply when plants are 4"-6" high (hooking stage). Subsequent applications should be at 2 to 3 week intervals.

SAFFLOWER, SUNFLOWER, AND OTHER OILSEED CROPS: Use 5 to 10 lbs. per acre per application.

SMALL GRAINS: (Wheat, oats, barley, rye) Use 5 to 10 lbs. per acre in sufficient water to cover by air or ground. Apply at tillering and again at early boot stage.

SUGAR BEETS: Use 5 to 10 lbs. per acre. Apply when leaves are 10" across. Repeat application 2 to 4 weeks later.

TOMATOES, PEPPERS: Use 5 to 10 lbs. per acre per application, beginning at pre-bloom. Repeat application every 2 to 3 weeks.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The directions for use of this product reflect the opinions of experts based on field uses and laboratory testing. Always follow label instructions. Manufacturer warrants that this product conforms to the product description on this label. Manufacturer or seller makes no other expressed or implied warranty. Neither manufacturer nor seller shall be liable for any injury or any application not specifically described on the label.

CAUTION:

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED - AVOID BREATHING DUST**

MANUFACTURED BY: JH BIOTECH, INC.
P.O. Box 3538
Ventura, CA 93006

NET WEIGHT: 50 LBS (22.5 Kg.)
 25 LBS (11.25 Kg.)