



BIOMIN[®]

BORON

2-0-0



Minerals for Soil and Foliar Application

PRODUCT INFORMATION

BIOMIN BORON Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct micronutrient deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizer.

BIOMIN BORON may be applied to **FRUIT TREES, FIELD CROPS, VEGETABLES, GREENHOUSE CROPS, POTATOES, GRAPES, CITRUS, ORNAMENTAL, GARDENS, etc...**

DIRECTION AND GENERAL RECOMMENDATIONS: SHAKE WELL BEFORE USING

SOIL APPLICATION:

BIOMIN BORON may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1 quart-1 gallon per acre (2.5-10 litres per hectare) depending on crop, soil conditions and application methods. For maintenance, 1 quart per acre (2.5 litres per hectare) is recommended.

FOLIAR APPLICATION:

FIELD CROPS: Apply 1-4 pints per acre (1.5-5 litres per hectare) of **BIOMIN BORON** when crops are 6-12 inches (15-30 cm) in height or when deficiency symptoms appear. Repeat application 2-4 weeks later.

VEGETABLES and POTATOES: Apply 1-4 pints per acre (1.5-5 litres per hectare) of **BIOMIN BORON** at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later. In greenhouses and plastic tunnels, application may be repeated every week or every other week.

GUARANTEED ANALYSIS

Boron (B) 3.0%

Derived from Boric Acid. Also contains Glycine (2% Nitrogen) and Citric Acid

PREVENT AND CORRECT MINERAL DEFICIENCIES

CAUTION:

This fertilizer contains Boron which may be injurious to certain crops. Contact your local county agency or field consultant for further information.

KEEP OUT OF REACH OF CHILDREN

Manufactured by:
JH BIOTECH, INC.
P.O. Box 7943
Ventura, California 93006

Information regarding the contents and levels of metals in this product is available on the internet at <http://agr.wa.gov>

FRUIT TREES AND GRAPES: Apply 1-4 pints per acre (1.5-5 litres per hectare) of **BIOMIN BORON** in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

For best results, apply **BIOMIN BORON** when plants are in an active growing stage, after irrigation or rainfall. For Foliar application, it should be diluted with at least 20 times of water. **BIOMIN BORON** will lower the high pH of water and decrease the hydrolysis of most pesticides.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Net Contents:

- 1 Pint (0.47 liter)
- 1 Gallon (3.78 liters)
- 2.5 Gallons (9.45 liters)
- 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters)
- 55 Gallons (207.9 liters)

Weight per Gallon: 9.7lbs. (4.4kg.)

LOT NO.:

0315-WA2

JH BIOTECH, INC.