

BIOMIN[®] MOLYBDENUM



Amino Acid Chelated Minerals for Soil and Foliar Application

PRODUCT INFORMATION

BIOMIN MOLYBDENUM - Amino acid chelated mineral for soil and/or foliar applications. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct Molybdenum deficiency and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

Biomin Molybdenum May Be Applied To Fruit Trees, Field Crops, Greenhouse Crops, Vegetables, Potatoes, Grapes, Citrus, Ornamental Gardens, Etc...

DIRECTIONS AND

GENERAL RECOMMENDATIONS:

SOIL APPLICATION

BIOMIN MOLYBDENUM may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1-5 quarts/acre (5-15 liters/hectare) depending on crop, soil conditions and application methods. For maintenance, 2 quarts per acre are usually recommended.

FOLIAR APPLICATION

FIELD CROPS: Apply 1-4 pints (2-4 liters/hectare) of **BIOMIN MOLYBDENUM** per acre when crops are 6-12 inches (15-30 cms) high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

VEGETABLES and POTATOES: Apply 1-4 pints (2-4 liters/hectare) of **BIOMIN MOLYBDENUM**

GUARANTEED ANALYSIS

Total Nitrogen (N)2.0%
2.0% Water Soluble Organic Nitrogen
Molybdenum (Mo).....3.0%
3.0% Chelated Molybdenum
Derived from Sodium Molybdate,
Citric Acid, and Glycine. Nitrogen
derived from Glycine.

PREVENTS AND CORRECTS MOLYBDENUM DEFICIENCY

KEEP OUT OF REACH OF CHILDREN

Manufactured by:

JH BIOTECH, INC.

P.O. Box 3538, Ventura, CA 93006

NET CONTENTS:

- 1 Pint
- 2.5 Gallons (9.45 liters)
- 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters)
- 55 Gallons (207.9 liters)
- Other: _____ Gallons

Weight per Gallon: 10.8 lbs. (4.9 kg.)

U.S. Pat.# 5,504,055

Lot No.:

per acre at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later.

GREENHOUSE CROPS: Apply 1-4 pints (2-4 liters/hectare) of **BIOMIN MOLYBDENUM** per acre 2-3 weeks after emergence or transplanting, or when **MOLYBDENUM** deficiencies first appear. Repeat the application every one or two weeks.

FRUIT TREES AND GRAPES: Apply 1-4 pints (2-4 liters/hectare) of **BIOMIN MOLYBDENUM** per acre in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

BIOMIN MOLYBDENUM is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water. For best results and more specific recommendations, contact your nearest distributor.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available online at <http://www.aapfco.org/metals.htm>

JH Biotech, Inc.

0305-D-A-01
Copyright © 2008 JH Biotech, Inc.
All rights reserved.